




CBGP
CENTRO DE BIOTECNOLOGÍA
Y GENÉTICA DE PLANTAS


Auxin - oxylipin crosstalk in stress responses of plant roots

CONGRESS OF THE
SOCIEDAD ESPAÑOLA DE BIOQUÍMICA Y BIOLOGÍA MOLECULAR
Barcelona, 07. September 2011


STEPHAN POLLMANN



The *aos/opr3* double mutant is a valuable tool for the uncoupled analysis of OPDA- and JA-triggered responses



The *aos/opr3* double mutant is a valuable tool for the uncoupled analysis of OPDA- and JA-triggered responses




Indole-3-acetic acid and Jasmonic acid - two on the first sight antagonistically acting signaling molecules




Auxins

- Growth promoting
- Apical dominance
- Tropisms
- Shoot elongation
- Induction of cambial cell division activity
- Formation of lateral- and adventitious roots
- Fruit development






Jasmonates

- Growth inhibiting
- Senescence promoting
- Plant defense
- Male fertility
- Flower development
- Mechanotransduction
- Tuber formation




Pollmann et al., *Plant Biol.* (2006); Böttcher & Pollmann, *FEBS J.* (2009); Lehmann et al., *Eur. J. Cell Biol.* (2010)

YUCCA flavin-containing monooxygenase-like proteins are recognized as key enzymes in auxin synthesis, catalyzing a rate-limiting reaction step




Photos courtesy of Youfa Cheng and Yunde Zhao

YUCCA flavin-containing monooxygenase-like proteins are recognized as key enzymes in auxin synthesis, catalyzing a rate-limiting reaction step




Photos courtesy of Youfa Cheng and Yunde Zhao


Transient expression of 35S::YUC8 and 35S::YUC9 after introduction into *Nicotiana benthamiana* leaves







Constitutive and stable overexpression of 35S::YUC8 and 35S::YUC9 in *Arabidopsis thaliana*




YUC8- and YUC9-GFP fusion proteins are exclusively located to the cytoplasm







qRT-PCR analyses corroborate the differential impact of oxylipins on *YUC8* and *YUC9* gene expression




yuc8- and *yuc9*-mutants can be distinguished from wild type as they have shorter hypocotyls and petioles as well as a reduced lateral root density





Relative to wild type, *yuc8*- and *yuc9*-mutants have reduced IAA contents and display reduced IAA production in response to MeJA treatment



yuc8- and *yuc9*-mutants are characterized by a reduced responsiveness to MeJA treatment





Acknowledgments

Current and former Group members:

Maik Hoffmann

Mathias Hentrich

Thomas Lehmann

Daniel Neu

Petra DÜchting

Collaborators:

Yunde Zhao, UCSD, USA

Hiroyuki Kasahara, RIKEN, Japan

Henrik Aronsson, GU, Sweden

Youfa Cheng, CAS, China

Jutta Ludwig-Müller, TUD, Germany

Josette Masle, ANU, Australia

Oliver Berkowitz, MurdochU, Australia

Markus Geisler, Unifr, Switzerland

Ronald Pierik, UU, Netherlands

Jesus Vicente Carbajosa, CBGP, Spain

Julia Kehr, CBGP, Spain

Juan Carlos del Pozo, CBGP, Spain

Stefano Boccaletti, CTB, Spain

Logos: CBGP (CENTRO DE BIOTECNOLOGÍA Y GENÓMICA DE PLANTAS), POLITÉCNICA, INIA (Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria), campus de Excelencia Internacional, Deutsche Forschungsgemeinschaft (DFG), SFB 480, MINISTERIO DE CIENCIA E INNOVACIÓN.